

hep-ph/0306219

CERN-TH/2003-138

UMN-TH-2205/03

FTPI-MINN-03/16

Updated Post-WMAP Benchmarks for Supersymmetry

M. Battaglia¹, A. De Roeck¹, J. Ellis¹, F. Gianotti¹, K. A. Olive² and L. Pape¹

¹*CERN, Geneva, Switzerland*

²*William I. Fine Theoretical Physics Institute,
University of Minnesota, Minneapolis, MN 55455, USA*

Abstract

We update a previously-proposed set of supersymmetric benchmark scenarios, taking into account the precise constraints on the cold dark matter density obtained by combining WMAP and other cosmological data, as well as the LEP and $b \rightarrow s\gamma$ constraints. We assume that R parity is conserved and work within the constrained MSSM (CMSSM) with universal soft supersymmetry-breaking scalar and gaugino masses m_0 and $m_{1/2}$. In most cases, the relic density calculated for the previous benchmarks may be brought within the WMAP range by reducing slightly m_0 , but in two cases more substantial changes in m_0 and $m_{1/2}$ are made. Since the WMAP constraint reduces the effective dimensionality of the CMSSM parameter space, one may study phenomenology along ‘WMAP lines’ in the $(m_{1/2}, m_0)$ plane that have acceptable amounts of dark matter. We discuss the production, decays and detectability of sparticles along these lines, at the LHC and at linear e^+e^- colliders in the sub- and multi-TeV ranges, stressing the complementarity of hadron and lepton colliders, and with particular emphasis on the neutralino sector. Finally, we preview the accuracy with which one might be able to predict the density of supersymmetric cold dark matter using collider measurements.

CERN-TH/2003-138

June 2003

1 Introduction

One of the crucial topics in the planning of analyses of data from experiments at present and future colliders is the search for supersymmetry [1]. Even the minimal supersymmetric extension of the Standard Model (MSSM), which conserves R parity, has over 100 free parameters once arbitrary soft supersymmetry-breaking parameters are allowed. For this reason, much attention is focussed on the constrained MSSM (CMSSM), in which the soft supersymmetry-breaking scalar masses m_0 , gaugino masses $m_{1/2}$ and trilinear parameters A_0 are each assumed to be universal at some high input scale, as in minimal supergravity and other models. These CMSSM parameters are constrained principally by the absence at LEP of new particles with masses $\lesssim 100$ GeV, by the agreement of $b \rightarrow s\gamma$ decay with Standard Model predictions, by the measurement of the anomalous magnetic moment of the muon, $g_\mu - 2$, and by the range allowed for relic cold dark matter, Ω_{CDM} .

Benchmark supersymmetric scenarios have a venerable history [2]. A couple of years ago, a new set of benchmark supersymmetric models was proposed in [3], consistent with all the above experimental constraints, as well as the cosmological constraint on Ω_{CDM} . Subsequently, there has not been any significant change in the LEP limits on supersymmetric particles [4, 5], we are still looking forward to more sensitive searches at the Fermilab Tevatron collider, the situation of $b \rightarrow s\gamma$ decay has changed little [6], and the interpretation of the $g_\mu - 2$ measurements remains unclear [7]. Various other benchmark scenarios have been proposed, in particular some supplementary points and lines in the CMSSM and in models with different mechanisms for supersymmetry breaking [8], benchmarks for supersymmetric Higgs physics [9], and scenarios in which the prospects for the Fermilab Tevatron collider are more favourable [10, 11].

The road(w)map for supersymmetric phenomenology has, however, been altered significantly by the recent improved determination of the allowable range of the cold dark matter density obtained by combining WMAP and other cosmological data: $0.094 < \Omega_{CDM} < 0.129$ at the $2\text{-}\sigma$ level [12]. This range is consistent with earlier indications, but more precise. Within the MSSM with conserved R parity, if one assumes that most of the cold dark matter consists of the lightest supersymmetric particle (LSP) and identifies this with the lightest neutralino χ [13], this WMAP constraint reduces the dimensionality of the parameter space. In the CMSSM, in particular, whereas generic regions of the $(m_{1/2}, m_0)$ planes would previously have been allowed for fixed values of $\tan\beta$ and A_0 [14], now only thin strips are permitted [15, 16] ¹.

¹Some authors had also considered an analogous narrow range for $\Omega_\chi h^2$ before this was mandated by the

It is of course possible to relax the restrictions to these narrow ‘WMAP lines’ in various ways. One could consider models that violate R parity, though these would also be subject to (different) astrophysical and cosmological constraints. Or one could consider alternatives to gravity-mediated supersymmetry breaking, such as anomaly- or gauge- mediated supersymmetry breaking, in which the sparticle spectra could be radically different, and/or the LSP might be a gravitino [8]. However, we do not discuss such models in this paper.

Here, we consider first the minor modifications of the previous CMSSM benchmark scenarios that bring them on to the WMAP lines. In most cases, this requires only a small change in m_0 , keeping $m_{1/2}$ the same as before, as seen in Table 1. An exception is the previous benchmark H, which had been chosen at the end of one of the coannihilation ‘tails’, at the largest possible value of $m_{1/2}$. The WMAP constraint not only reduces the possible spread in m_0 for fixed $m_{1/2}$, but also reduces the allowable range of $m_{1/2}$. Therefore, our modification of benchmark H has reduced values of both $m_{1/2}$ and m_0 . Another example is benchmark M, which has also been shifted significantly because the rapid-annihilation ‘funnel’ for $\tan\beta = 50$ and $\mu > 0$ is affected substantially by the WMAP restriction on $\Omega_\chi h^2$. More detailed discussions of these updated benchmark points are given in Section 2, where we also discuss the (generally small) extent to which the decay branching ratios differ from the previous versions of the benchmark points ².

Then, in Section 3, we discuss systematically how the phenomenology of supersymmetric models varies with $m_{1/2}$ along the WMAP lines. We first present parametrizations of these lines appropriate for the SSARD [18] and ISASUGRA 7.67 [20] codes. We then discuss the patterns of decays of various selected sparticles. We show in particular that $\chi_2 \rightarrow \tilde{\tau}_1 \tau$ decays are generally important, though other $\chi_2 \rightarrow \tilde{\ell} \ell$ decays (where $\ell \equiv e, \mu$) are also important for low $m_{1/2}$ at low $\tan\beta$. The decays $\chi_2 \rightarrow \chi h, \chi Z$ are never very large on the WMAP lines, despite being kinematically allowed for most values of $m_{1/2}$. We then discuss the average numbers of Z, h and τ particles produced per sparticle production event at the LHC. This study confirms the importance of the τ signature, with trilepton signatures also looking promising. We discuss the extent to which our various benchmark scenarios sample generic features along the WMAP lines.

Section 4 contains a discussion of sparticle detectability at various accelerators, starting with the LHC ³. We find that the lightest MSSM Higgs boson and all the squarks are in prin-

WMAP data [17].

²Where it seems necessary to avoid confusion with the previous versions of these benchmark scenarios [3], we denote the updated versions with primes: A’, B’, etc.: otherwise we retain the same notation.

³We have nothing to add here to the discussion in [3] of the prospects for detecting supersymmetry at the Fermilab Tevatron collider, referring the reader to [10, 11] for alternative discussions.

Supersymmetric spectra in post-WMAP benchmark scenarios

Model	A'	B'	C'	D'	E'	F'	G'	H'	I'	J'	K'	L'	M'
$m_{1/2}$	600	250	400	525	300	1000	375	935	350	750	1300	450	1840
m_0	120	60	85	110	1530	3450	115	245	175	285	1000	300	1100
$\tan \beta$	⁽¹⁴⁰⁾ 5	⁽¹⁰⁰⁾ 10	⁽⁹⁰⁾ 10	⁽¹²⁵⁾ 10	⁽¹⁵⁰⁰⁾ 10	10	⁽¹²⁰⁾ 20	⁽⁴¹⁹⁾ 20	⁽¹⁸⁰⁾ 35	⁽³⁰⁰⁾ 35	35	⁽³⁵⁰⁾ 50	⁽¹⁵⁰⁰⁾ 50
$\text{sign}(\mu)$	+	+	+	−	+	+	+	+	+	+	−	+	+
$\alpha_s(m_Z)$	121	125	123	121	124	120	124	120	123	120	118	122	117
m_t	175	175	175	175	171	171	175	175	175	175	175	175	175
Masses													
$ \mu(m_Z) $	741	333	503	634	205	496	471	1026	439	843	1317	540	1764
h	115	113	116	117	114	118	117	122	116	121	118	118	124
H	884	375	578	736	1532	3491	523	1185	452	883	1176	489	1652
A	882	375	578	735	1533	3491	523	1185	451	883	1176	489	1663
H $^\pm$	886	383	584	740	1535	3492	529	1188	459	887	1180	496	1654
χ_1	252	98	163	220	115	430	153	402	143	320	573	187	821
χ_2	480	181	310	424	182	522	289	774	270	615	1105	358	1583
χ_3	761	346	519	655	221	523	489	1068	464	897	1413	588	1994
χ_4	775	365	535	662	304	885	504	1078	478	906	1421	599	1999
$\chi_1^\pm$	480	180	309	424	174	511	290	774	270	615	1105	358	1583
$\chi_2^\pm$	775	367	535	664	304	886	505	1079	479	907	1422	600	1999
$\tilde{g}$	1715	715	1145	1495	869	2914	1075	2681	999	1593	3716	994	5262
e_L, μ_L	425	188	289	375	1544	3512	285	673	300	581	1319	430	1635
e_R, μ_R	261	121	180	232	1535	3471	189	433	224	405	1114	348	1300
ν_e, ν_μ	418	171	278	367	1542	3511	273	669	289	575	1317	422	1633
τ_1	258	112	172	225	1522	3443	162	403	155	323	971	200	920
τ_2	425	192	291	376	1538	3498	291	670	310	573	1268	420	1511
ν_τ	418	187	277	366	1542	3497	270	661	277	555	1261	386	1502
u_L, c_L	1202	546	834	1064	1644	3908	792	1808	755	1493	2602	965	3491
u_R, c_R	1151	527	803	1021	1635	3867	762	1730	723	1429	2494	930	3332
d_L, s_L	1205	552	838	1067	1646	3909	797	1810	758	1429	2603	968	3492
d_R, s_R	1144	526	799	1016	1634	3861	759	1718	723	1495	2479	925	3309
t_1	896	393	618	807	1050	2580	587	1380	553	1131	1935	710	2630
t_2	1143	573	819	1013	1387	3330	777	1677	731	1372	2237	891	3054
b_1	1101	502	765	976	1379	3323	717	1645	659	1325	2173	815	2998
b_2	1144	528	798	1011	1622	3834	756	1695	711	1377	2242	880	3062

Table 1:

Proposed post-WMAP CMSSM benchmark points and mass spectra (in GeV), as calculated using SSARD [18] and FeynHiggs [19], using the one-loop corrected effective potential computed at the electroweak scale and one-loop corrections to the chargino and neutralino masses. We recall (in parentheses) the values of $m_{1/2}$ and m_0 used in [3], in cases where they differ. As in [3], exact gauge coupling unification is enforced, resulting in the predictions for $\alpha_s(m_Z)$ that are shown in units of 0.001. We use $A_0 = 0$, $m_b(m_b)^{\overline{MS}} = 4.25$ GeV and $m_t = 175$ GeV for most of the points, with the exceptions of points E' and F', where $m_t = 171$ GeV is used.

ciple observable along the complete WMAP lines, except along parts of the rapid-annihilation ‘funnels’, whereas sleptons and other neutralinos and charginos are detectable only in the lower parts of the WMAP ranges of $m_{1/2}$. This discussion is extended in Subsection 4.2 to linear e^+e^- colliders with $E_{CM} = 0.5, 1.0$ [21] and 3.0 or 5.0 TeV [22]. Confirming previous studies [21, 3, 8], we see that a machine with $E_{CM} = 0.5$ TeV would be able to explore supersymmetry in the lower parts of the WMAP ranges of $m_{1/2}$, whereas a machine with $E_{CM} = 1.0$ TeV would be able to explore supersymmetry along (almost) the entire WMAP lines. CLIC with $E_{CM} = 3.0$ TeV would be able to complete the spectrum of electroweakly-interacting sparticles along the entire WMAP lines, as well as furnish detailed measurements of squarks for a large fraction of the allowed range of $m_{1/2}$ [23]. We also discuss in more detail the observability of neutralinos at the different colliders surveyed, with particular emphasis on CLIC studies.

Finally, in Section 5 we review some conclusions from our results, addressing in particular the question whether collider measurements will have the potential to determine if the LSP constitutes most of the cold dark matter in the Universe [13, 24].

2 Updated CMSSM Benchmark Points

2.1 Improved Choices of Supersymmetry-Breaking Parameters

Since the laboratory constraints on the CMSSM have not yet changed substantially since the termination of LEP [4], the only reason for updating the benchmark points we proposed previously [3] is the refined estimate of $\Omega_{CDM}h^2$ provided by combining the new WMAP data with those previously available [12]. Assuming that most of the cold dark matter is composed of LSPs χ , previously we allowed $0.1 < \Omega_\chi h^2 < 0.3$ (rather conservatively), whereas the WMAP analysis now allows only the range

$$0.094 < \Omega_\chi h^2 < 0.129 \quad (1)$$

at the $2\text{-}\sigma$ level [12]. We see in Table 2 of [3] that *all* the previous benchmark points yielded relic densities $\Omega_\chi h^2$ *above* the range (1). Since the relic density calculations have not changed significantly since [3] in the regions of interest, the previous benchmark points must be moved.

Of the 13 benchmark points proposed previously, 5 (B, C, G, I, L) were in the ‘bulk’ regions at low $m_{1/2}$ and m_0 , 4 (A, D, H, J) were along the coannihilation ‘tails’ extending to larger $m_{1/2}$ [25], 2 (K, M) were along rapid-annihilation ‘funnels’ where both $m_{1/2}$ and m_0 may grow large [26], and 2 (E, F) were in the ‘focus-point’ region at very large m_0 [27].

The WMAP constraint (1) has slimmed the ‘bulk’ region down considerably, but only minor reductions in m_0 are needed to reduce the relic density of points (B, C, G, I, L) into the range (1), as shown in Table 1. The coannihilation ‘tails’ are also much thinner than they were before, and the required reductions in m_0 for points (A, D, J) are also small. The previous point H is an exception to this rule, since it was chosen at the extreme tip of a coannihilation ‘tail’. In this case, since the upper limit in (1) is considerably reduced compared with that assumed in [3], the allowed upper limit on $m_{1/2}$ has been reduced substantially [15]. Hence we must make reductions in both $m_{1/2}$ and m_0 for point H, as also shown in Table 1. In the case of point K, the rapid-annihilation ‘funnel’ has become thinner following WMAP, and a minor adjustment in $m_{1/2}$ is sufficient, but in the case of point M, the ‘funnel’ has changed substantially, and both $m_{1/2}$ and m_0 had to be changed.

The two focus-point benchmarks (E, F) may also be adapted to the WMAP constraint (1) with small changes, as also shown in Table 1, which is based on the **SSARD** code. However, we take this opportunity to underline the extreme delicacy of model calculations in this region, as may be seen by comparing the results of different codes [18, 20, 28, 29] in their successive releases ⁴. The various codes do agree that there is a strip in the ‘focus-point’ region where $\Omega_\chi h^2$ is in the WMAP range (1), but the error in its location due to the experimental uncertainty in the mass of the top quark m_t (for example) is much larger than the intrinsic width of the WMAP strip in this region. We have chosen a pole mass $m_t = 171$ GeV for benchmarks E and F, as opposed to the choice $m_t = 175$ GeV made for the other benchmarks, which would have required much larger values of m_0 in this focus-point region, namely $m_0 = 2550, 5030$ GeV for the **SSARD** code, respectively. The new D0 value of $m_t \simeq 179$ GeV [30], if confirmed, would push the ‘focus-point’ region up to still larger m_0 , whatever code is used: specifically to $m_0 = 5800, 9070$ GeV for benchmarks E and F, respectively, if the **SSARD** code is used ⁵.

There are also significant theoretical uncertainties associated with higher-order effects [31], that are reflected in differences between different codes in their various versions. For example, for $m_t = 175$ GeV at benchmarks E and F (with $m_{1/2} = 300, 1000$ GeV, respectively), the **ISASUGRA 7.51**, [**ISASUGRA 7.67**], [**SUSPECT 2.10**], [**SUSPECT 2.11**] codes require $m_0 = 1530, 3450$ GeV, [$m_0 = 3590, 6260$ GeV], [$m_0 = 2350, 4110$ GeV], [$m_0 = 2590, 3850$ GeV], respectively. We see that the predictions of successive versions of the **SUSPECT** code [28] do not vary much, and agree quite well with **SSARD** [18], whereas there has been significant evolution in the predictions of **ISASUGRA** [20].

⁴See also the discussion in [3].

⁵This would put all sfermions beyond the reach of all the colliders discussed here.

In view of these uncertainties in the focus-point region, we concentrate in the following on the remaining updated benchmark points proposed in Table 1. These are located on the ‘WMAP lines’ shown in Fig. 1(a) for $\mu > 0$ and $\tan\beta = 5, 10, 20, 35$ and 50, and (b) for $\mu < 0$ and $\tan\beta = 10$ and 35⁶. We recall that, for given values of $\tan\beta$, $m_{1/2}$ and the sign of μ , lower values of m_0 generally have values of $\Omega_\chi h^2$ below the WMAP range⁷, whereas higher values of m_0 generally have values of $\Omega_\chi h^2$ that are too high, and are therefore unacceptable unless one goes beyond the CMSSM framework used here. At the ends of the ‘WMAP lines’, smaller values of $m_{1/2}$ are excluded by either m_h [5, 19] and/or $b \rightarrow s\gamma$ [6], and larger values of $m_{1/2}$ have values of $\Omega_\chi h^2$ above the WMAP range and/or the LSP is the charged $\tilde{\tau}_1$. We note that, as was commented in [3], most of the proposed benchmark points for $\mu > 0$ yield a value of $g_\mu - 2$ that lies within 2σ of the present experimental value based on e^+e^- data, corresponding to the lighter (pink) regions of the strips in Fig. 1(a). However, we do not impose this as a requirement on the benchmark points, as exemplified in Fig. 1(b) for $\mu < 0$.

We see also in Fig. 1 that the majority of the benchmark points lie on the portions of the WMAP lines with lower values of $m_{1/2}$, making them more interesting for the early stages of LHC operation, and for any sub-TeV linear e^+e^- collider. Generally speaking, less fine-tuning of parameters is required in this region than in the focus-point, funnel and tail regions [32]: see also the discussion in [3].

2.2 Discussion of Spectra

In order to study the detectability of MSSM particles and the possible accuracies in measurements for the proposed benchmarks, it is essential to relate the input CMSSM parameters, defined in Table 1 for the **SSARD** program, to those necessary to obtain an equivalent MSSM spectrum in Monte Carlo generators⁸. Accordingly, we have tuned the inputs for **ISASUGRA 7.67** to reproduce as accurately as possible the spectra given in Table 1, by scanning m_0 and $\tan\beta$ to identify the parameter pairs minimizing the sum of the relative differences of particle masses $|m_{\text{SSARD}} - m_{\text{ISASUSY}}|/M_{\text{SSARD}}$. In view of their importances for the calculation of $\Omega_\chi h^2$, we have treated specially the mass splitting $|m_{\tilde{\tau}_1} - m_\chi|$, which is important along the coannihilation ‘tail’, and also $|m_A - 2m_\chi|$, where it becomes important in the ‘funnel’ regions. We have given each of them 30 % of the total weighting when optimizing the **Isasugra 7.67** parameters in the relevant $(m_{1/2}, m_0)$ regions. The results of the optimization are given in

⁶There would be corresponding WMAP lines in the focus-point regions, whose location is uncertain, but which would generally lie at larger values of m_0 .

⁷Therefore, in this region the LSP might constitute only a part of the cold dark matter.

⁸See also the discussion in [3].




Figure 1: The shaded strips display the regions of the $(m_{1/2}, m_0)$ plane that are compatible [15] with $0.094 < \Omega_\chi h^2 < 0.129$ in the ‘bulk’, coannihilation ‘tail’, and rapid-annihilation ‘funnel’ regions, as well as the laboratory constraints, for (a) $\mu > 0$ and $\tan \beta = 5, 10, 20, 35$ and 50 , and (b) for $\mu < 0$ and $\tan \beta = 10$ and 35 . The parts of these ‘WMAP lines’ for $\mu > 0$ compatible with $g_\mu - 2$ at the 2- σ level have lighter (pink) shading [7]. The updated post-WMAP benchmark scenarios are marked in red. Points (E', F') in the focus-point region have larger values of m_0 .

Table 2.

In general, good matching between the predictions of the two codes can be found with only moderate shifts of the input parameters, and we have chosen only to adjust m_0 in most cases. The typical average relative difference in the benchmark point masses is of the order of a few percent. However, occasionally individual masses may exhibit discrepancies of up to 15%, and ensuring compatible mass splittings ΔM requires a systematic shift in m_0 to higher values, particularly at large $\tan\beta$, as seen in Fig. 2, where the cases $\tan\beta = 5, 10, 20, 35$ and 50 are exhibited for $\mu > 0$, and $\tan\beta = 10$ for $\mu < 0$. In addition to increasing m_0 , here we find a better correspondence between **SSARD** and **ISASUGRA 7.67** if $\tan\beta$ is allowed to be reduced, as done for benchmarks (K, L, M) in Table 2.

We have further studied whether the resulting **ISASUGRA 7.67** versions of the benchmark points yield values of $\Omega_\chi h^2$ that are indeed compatible with the WMAP data. The first line of Table 3 shows the values of $\Omega_\chi h^2$ calculated using the **SSARD** [18] code at the updated benchmark points shown in Table 1, which are well within the WMAP range. The second line shows the corresponding values of $\Omega_\chi h^2$ calculated using the **Micromegas** code [33] interfaced with the **ISASUGRA 7.67** code that is used to calculate the spectra in Table 2. The agreement between **SSARD** and **Micromegas/ISASUGRA 7.67** is generally good in the bulk and coannihilation tail regions, with the latter falling within the WMAP range, except for point H (which is right at the tip of the coannihilation tail) and point J (where the discrepancy is only marginal). The **Micromegas/ISASUGRA 7.67** code gives results that are outside the WMAP range in the focus-point and at the upper edge of the funnel region (points E, F and M), reflecting the fact that the calculation of $\Omega_\chi h^2$ is notoriously delicate in these regions [32] and that the tuning of the mass spectra between the two different codes becomes difficult.

For completeness, we also present in Table 2 the values of the $b \rightarrow s\gamma$ branching ratio and the supersymmetric contribution to $g_\mu - 2$ calculated using **SSARD**. These may be compared with any evolution in the future experimental values of these quantities. As already mentioned, the situation with regard to any possible non-Standard Model contribution to the experimental value of $g_\mu - 2$ is sufficiently volatile that we do not use this as a criterion for selecting benchmarks, though it is used in Fig. 3 to order the benchmark points, as described below.

**Supersymmetric spectra in post-WMAP benchmarks
calculated with ISASUGRA 7.67**

Model	A'	B'	C'	D'	E'	F'	G'	H'	I'	J'	K'	L'	M'
$m_{1/2}$	600	250	400	525	300	1000	375	935	350	750	1300	450	1840
m_0	107	57	80	101	1532	3440	113	244	181	299	1001	303	1125
$\tan\beta$	5	10	10	10	10	10	20	20	35	35	46	47	51
$\text{sign}(\mu)$	+	+	+	-	+	+	+	+	+	+	-	+	+
m_t	175	175	175	175	171	171	175	175	175	175	175	175	175
Masses													
$ \mu(m_Z) $	773	339	519	663	217	606	485	1092	452	891	1420	563	1940
h	116	113	117	117	114	118	117	122	117	121	123	118	124
H	896	376	584	750	1544	3525	525	1214	444	888	1161	480	1623
A	889	373	580	745	1534	3502	522	1206	441	882	1153	477	1613
$H^\pm$	899	384	589	754	1546	3524	532	1217	453	892	1164	490	1627
χ	242	95	158	212	112	421	148	388	138	309	554	181	794
χ_2	471	180	305	415	184	610	286	750	266	598	1064	351	1513
χ_3	778	345	525	671	229	622	492	1100	459	899	1430	568	1952
χ_4	792	366	540	678	302	858	507	1109	475	908	1437	582	1959
$\chi_1^\pm$	469	178	304	415	175	613	285	750	265	598	1064	350	1514
$\chi_2^\pm$	791	366	541	679	304	846	507	1108	475	908	1435	582	1956
$\tilde{g}$	1367	611	940	1208	800	2364	887	2061	835	1680	2820	1055	3884
e_L, μ_L	425	188	290	376	1543	3499	285	679	304	591	1324	434	1660
e_R, μ_R	251	117	174	224	1534	3454	185	426	227	410	1109	348	1312
ν_e, ν_μ	412	167	274	362	1539	3492	270	665	290	579	1315	423	1648
τ_1	249	109	167	217	1521	3427	157	391	150	312	896	194	796
τ_2	425	191	291	376	1534	3485	290	674	312	579	1251	420	1504
ν_τ	411	167	273	360	1532	3478	266	657	278	558	1239	387	1492
u_L, c_L	1248	558	859	1103	1639	3923	814	1885	778	1554	2722	1001	3670
u_R, c_R	1202	542	830	1064	1637	3897	787	1812	754	1497	2627	969	3528
d_L, s_L	1250	564	863	1107	1641	3924	818	1887	783	1556	2723	1004	3671
d_R, s_R	1197	541	828	1059	1638	3894	786	1804	752	1491	2615	967	3509
t_1	958	411	653	860	1052	2647	617	1477	584	1207	2095	753	2857
t_2	1184	576	837	1048	1387	3373	792	1753	748	1428	2366	920	3231
b_1	1147	514	789	1015	1375	3356	737	1719	677	1377	2297	844	3149
b_2	1181	535	816	1043	1602	3816	770	1761	725	1423	2349	904	3217

Table 2: *Proposed post-WMAP CMSSM benchmark points and mass spectra (in GeV), as calculated using ISASUGRA 7.67 [20] and adapting the values of m_0 and $\tan\beta$ (when it is large) to give the best fit to the SSARD spectra shown in Table 1, as described in the text.*




Figure 2: Comparison of the mass differences $\Delta M \equiv |m_{\tilde{\tau}_1} - m_\chi|$ along the WMAP lines, as calculated in the SSARD code (dots) and ISASUGRA 7.67 using the same values of m_0 (dashed lines), as functions of $m_{1/2}$ for $\mu > 0$ and $\tan \beta = 5, 10, 20, 35, 50$ and $\mu < 0, \tan \beta = 10$. We see the need to shift m_0 systematically, particularly at larger $m_{1/2}$ and $\tan \beta$. This has been done for the solid lines, which are improved ISASUGRA 7.67 fits to the WMAP lines described in the text.

Relic density $\Omega_\chi h^2$, $b \rightarrow s\gamma$ and $g_\mu - 2$ in post-WMAP benchmark scenarios

	A'	B'	C'	D'	E'	F'	G'	H'	I'	J'	K'	L'	M'
$\Omega_\chi h^2$: SSARD	.12	.12	.12	.10	.10	.10	.13	.13	.12	.10	.10	.10	.14
$\Omega_\chi h^2$: Micromegas	.09	.12	.12	.09	.33	2.56	.12	.16	.12	.08	0.12	.11	0.27
$\text{BR}(b \rightarrow s\gamma) \times 10^4$	3.9	3.1	3.8	4.5	3.7	3.7	3.3	3.6	2.5	3.4	4.2	2.5	3.5
$\delta a_\mu \times 10^9$	0.3	3.2	1.3	-0.8	0.2	0.03	2.7	0.4	4.5	1.1	-0.3	3.4	0.3

Table 3: Comparison of $\Omega_\chi h^2$ for the benchmark points in Table 1 computed with the SSARD code [18] (top line), and the Micromegas code [33] interfaced with ISASUGRA 7.67 [20] (second line) using the fitted parameters shown in Table 2. The third and fourth lines show the values of $b \rightarrow s\gamma$ and $g_\mu - 2$ calculated using SSARD.

2.3 Observability at Different Accelerators

As an important ingredient in the subsequent analyses of sparticle observability at different colliders, we first report a comparison of the principal sparticle branching ratios at the updated benchmark points with those found previously [3] at the original benchmark points. The new branching ratios have been calculated using ISASUGRA 7.67, whereas previously they were calculated using ISASUGRA 7.51 [20], and we comment on the differences associated with the improvement in ISASUGRA, as opposed to the changes in the benchmark parameter choices⁹. Since the differences in branching ratios are not large, in general, we limit ourselves to commenting on instances where the more significant differences occur.

- In the case of point A, the most notable difference is an increase in $B(\tilde{b}_2 \rightarrow b\chi)$ from 39 to 62 %, with corresponding decreases in $B(\tilde{b}_2 \rightarrow t + \chi^-, \tilde{t}_1 + W^-)$. There is also an increase of $\chi_4 \rightarrow h\chi_2$ from 23 to 29 % at the expense of $\chi_4 \rightarrow W\chi_1^\pm$. These changes are not related to the new parameter choice, but to the differences in the ISASUGRA versions.

- In the case of point B, there is again an increase in $B(\tilde{b}_2 \rightarrow b\chi)$, from 15 to 25 %, and a decrease in $B(\tilde{b}_2 \rightarrow \tilde{t}_1 + W^-)$. There is also an increase in $B(\tilde{e}_L \rightarrow e\chi)$ from 47 to 87 %, accompanied by decreases in $B(\tilde{e}_L \rightarrow e\chi_2, \nu + \chi_1^\pm)$. We also note that the $B(\tilde{\nu}_\ell \rightarrow \ell^- \chi^+)$ now vanish, so that the $\tilde{\nu}_\ell$ are *invisible*. There is also an increase in $B(\tilde{\tau}_2 \rightarrow \tau\chi)$ from 51 to 84 %, which decreases $B(\tilde{\tau}_2 \rightarrow \nu_\tau + \chi^-, \tau + \chi_2)$. Finally, there are decreases in $B(\chi_2 \rightarrow \tilde{\tau}_1 + \tau)$ from 84 to 42 %, which increases the branching ratio for the invisible channel $B(\chi_2 \rightarrow \tilde{\nu} + \nu)$, and in $B(\chi^\pm \rightarrow \tilde{\tau}_1 + \nu_\tau)$ from 96 to 36 %, which increases $B(\chi^\pm \rightarrow \tilde{\nu} + \ell)$. The changes for the sleptons are mostly because they are lighter at the updated benchmark point with smaller m_0 , and those for $\tilde{b}_2$ are related to the differences in the ISASUGRA versions. Both effects

⁹For the purposes of our discussion, one of the most important upgrades to ISASUGRA has been the inclusion of one-loop radiative corrections to sparticle masses.

influence the gaugino branching ratios.

- At benchmarks C and D, $B(\tilde{b}_2 \rightarrow b\chi)$ increases from 13 to 24 % and from 20 to 44 %, respectively, with corresponding decreases in $B(\tilde{b}_2 \rightarrow \tilde{t}_1 W^-)$ and $B(\tilde{b}_2 \rightarrow t + \chi^\pm)$. At point C, decreases are observed of $\chi_2 \rightarrow \tilde{\tau}_1 \tau$ from 23 to 13 % and of $\chi_1^\pm \rightarrow \tilde{\tau}_1 \nu$ from 21 to 11 %, compensated by increases of $\chi_2 \rightarrow \tilde{\nu} \nu$ and $\chi_1^\pm \rightarrow \tilde{\nu} l$, respectively. All changes are mostly related to differences in the ISASUGRA versions.

- At the updated points E and F, the most notable effect is an increase of the branching ratios of squarks to gluinos, due to the increase in m_0 and hence of the squark masses at the updated benchmark points. For instance, the $B(\tilde{b}_1 \rightarrow b\tilde{g})$ changes from 45 to 52 % at E and from 22 to 38 % at F. They are compensated by a reduction of the decays to charginos and neutralinos. Moreover, at point E the decays of sleptons into $\chi_1^\pm$ tend to decrease and those into $\chi_2^\pm$ to increase correspondingly.

- At the updated benchmark G, there are increases in $B(\tilde{b}_1 \rightarrow t\chi_2^\pm)$, from 13 to 22 %, with accompanying decreases in other modes, mainly in $B(\tilde{b}_1 \rightarrow t + \chi_1^\pm, b + \chi_2)$. A decrease is observed of $\chi_2 \rightarrow \tilde{\tau}_1 \tau$ from 82 to 62 % and of $\chi_1^\pm \rightarrow \tilde{\tau}_1 \nu$ from 81 to 57 %, compensated by an increase of $\chi_2 \rightarrow \tilde{\nu} \nu$ and $\chi_1^\pm \rightarrow \tilde{\nu} l$, respectively. The changes are mostly due to the differences in the ISASUGRA versions.

- Although the updated point H has a significantly lighter spectrum than previously, no major changes (> 5 -6 %) are observed in the decay branching ratios.

- For the old point I the decay $\tilde{b}_1 \rightarrow t\chi_2^\pm$ was kinematically forbidden. For the updated point it is allowed and has a branching ratio of 11 %. Also, $B(\tilde{\nu}_e \rightarrow \nu\chi)$ increases from 61 to 73 % and $B(\tilde{\nu}_e \rightarrow e\chi_1^\pm)$ decreases correspondingly, mostly due to the change in the ISASUGRA version used.

- At point J the main changes are a decrease of $B(\chi_2 \rightarrow \tilde{\tau}_1 \tau)$ from 82 to 66 % and of $B(\chi_1^\pm \rightarrow \tilde{\tau}_1 \nu)$ from 82 to 64 %, mostly compensated by an increase of $B(\chi_2 \rightarrow \tilde{\nu} \nu)$ and $B(\chi_1^\pm \rightarrow \tilde{\nu} l)$, respectively. The changes are mainly due to the differences in the ISASUGRA versions.

- At the updated benchmark K, there is an increased branching ratio for the direct decay of $\tilde{e}_L$ and $\tilde{\nu}_e$ to χ from 26 to 35 % and 27 to 37 % respectively, at the expense of the decays to $\chi_1^\pm$. The increase of $B(\tilde{\nu}_\tau \rightarrow W\tilde{\tau}_1)$ from 28 to 40 % is accompanied by a decrease of $B(\tilde{\nu}_\tau \rightarrow \tau\chi_1^\pm)$. Also, the decays of χ_2 and $\chi_1^\pm$ to $\tilde{\tau}_1$, which were previously forbidden kinematically, are now allowed by the new parameter values and reach 43 and 41 %, respectively. A reduction of $B(\chi_4 \rightarrow W\chi_1^\pm)$ from 56 to 47 % is also observed, associated with the differences in the ISASUGRA versions.

- The updated point L is characterized by a decrease of the branching ratios of $\tilde{\nu}_e$ and $\tilde{\nu}_\tau$

to the $\chi_1^\pm$, compensated by an increase of $B(\tilde{\nu}_e \rightarrow \nu\chi)$ from 35 to 43 % and of $B(\tilde{\nu}_\tau \rightarrow W\tilde{\tau}_1)$ from 59 to 77 %. The decay branching ratios of χ_3 and χ_4 to $\tilde{\tau}_1\tau$ are increased from 19 to 27 % and 13 to 20 %, respectively, with a corresponding reduction of the decays to $\chi_1^\pm$. These changes are associated with the updated CMSSM parameter values.

- At the updated point M, the main change is a smaller value of m_0 , which affects many branching fractions. We only mention some of the major changes. The direct decay to χ is increased for $\tilde{e}_L$ and $\tilde{\nu}_e$ from 34 to 74 % and from 34 to 77 %, respectively, at the expense of a reduced decay to the chargino and the heavier neutralino. The $B(\chi_2 \rightarrow h + \chi)$ decreases from 45 to 12 %, whereas $B(\chi_2 \rightarrow \tilde{\tau}_1 + \tau)$ increases, and $B(\chi_1^\pm \rightarrow W^\pm + \chi)$ decreases from 48 to 14 %, whereas $B(\chi_1^\pm \rightarrow \tilde{\tau}_1 + \nu_\tau)$ increases.

The branching ratios at the updated benchmark points not mentioned above are not significantly different from those at the original versions of the points. Among all the above effects, perhaps the most significant is the invisibility of the $\tilde{\nu}_\ell$ at the updated version of point B.

Combining the above information, we now present in Fig. 3 an updated comparison of the numbers of different MSSM particles that should be observable at different accelerators in the various benchmark scenarios [3], ordered by their consistency with $g_\mu - 2$ as calculated using e^+e^- data for the Standard Model contribution [7]. We re-emphasize that the qualities of the prospective sparticle observations at hadron colliders and linear e^+e^- colliders are often very different, with the latter's clean experimental environments providing prospects for measurements with better precision. Nevertheless, Fig. 3 already restates the clear message that hadron colliders and linear e^+e^- colliders are largely complementary in the classes of particles that they can see, with the former offering good prospects for strongly-interacting sparticles such as squarks and gluinos, and the latter excelling for weakly-interacting sparticles such as charginos, neutralinos and sleptons. We discuss later the detailed criteria used for assessing the detectabilities of different particles at different colliders.

3 WMAP Lines

As has already been mentioned, in view of the reduction in dimensionality of the CMSSM parameter space enforced by WMAP [15], one may progress beyond the previous approach of sampling, more or less sparsely, the CMSSM parameter space. In particular, as proposed in [8] but with a different attitude towards the cosmological density constraint, one may explore, more or less systematically, the CMSSM phenomenology along the lines in parameter space shown in Fig. 1. Again as proposed in [8], one may then focus special attention




Figure 3: Summary of the numbers of MSSM particles that may be detectable at various accelerators in the updated benchmark scenarios. As in [3], we see that the capabilities of the LHC and of linear e^+e^- colliders are largely complementary. We re-emphasize that mass and coupling measurements at e^+e^- colliders are usually much cleaner and more precise than at hadron-hadron colliders such as the LHC, where, for example, it is not known how to distinguish the light squark flavours.

on particular points along these lines. The updated benchmark points discussed in the previous section are examples and, as we discuss below, they manifest some of the distinctive possibilities that may appear along the WMAP lines. However, for certain purposes it may be helpful to select a few additional points that exemplify other generic possibilities, as we discuss later.

3.1 Parametrizations of WMAP Lines

As seen in Fig. 1, the WMAP lines have a relatively simple form for $\tan\beta \lesssim 35$. It is therefore possible to provide simple quadratic parametrizations for them that are a convenient summaries of the correlations between $m_{1/2}$ and m_0 that are enforced by WMAP within the CMSSM framework adopted here. Using the `SSARD` code, we find the following convenient parametrizations for the $\mu > 0$ lines:

$$\tan\beta = 5 : \quad m_0 = -5.46 + 0.206m_{1/2} + 4.02 \times 10^{-6}m_{1/2}^2, \quad (475 \leq m_{1/2} \leq 960) \quad (2)$$

$$\tan\beta = 10 : \quad m_0 = 11.97 + 0.163m_{1/2} + 4.90 \times 10^{-5}m_{1/2}^2, \quad (300 \leq m_{1/2} \leq 940) \quad (3)$$

$$\tan\beta = 20 : \quad m_0 = 50.89 + 0.144m_{1/2} + 6.77 \times 10^{-5}m_{1/2}^2, \quad (270 \leq m_{1/2} \leq 940), \quad (4)$$

$$\tan\beta = 35 : \quad m_0 = 90.24 + 0.219m_{1/2} + 5.70 \times 10^{-5}m_{1/2}^2, \quad (315 \leq m_{1/2} \leq 1070) \quad (5)$$

where the masses are expressed in units of GeV, and the numbers in parentheses specify the allowed ranges of $m_{1/2}$. When $\tan\beta = 50$, a rapid-annihilation funnel appears, and parametrizing the regions allowed by WMAP becomes more complicated. In the case of $\tan\beta = 50$, one may conveniently use:

$$\begin{aligned} m_0 &= 140.6 + 0.33m_{1/2} + 6.3 \times 10^{-5}m_{1/2}^2 \quad (350 \leq m_{1/2} \leq 1550), \\ m_0 &= -1941 + 1.65m_{1/2} \quad (1700 \leq m_{1/2} \leq 1800), \end{aligned} \quad (6)$$

where the masses are again expressed in GeV units. For $\mu < 0$, we propose the following convenient parametrizations:

$$\tan\beta = 10 : \quad m_0 = 15.0 + 0.16m_{1/2} + 4.2 \times 10^{-5}m_{1/2}^2, \quad (400 \leq m_{1/2} \leq 940) \quad (7)$$

$$\tan\beta = 35 : \quad m_0 = 39.6 + 0.76m_{1/2} - 1.9 \times 10^{-5}m_{1/2}^2, \quad (790 \leq m_{1/2} \leq 1650), \quad (8)$$

where for $\tan\beta = 35$ we parametrize only the top branch allowed by WMAP in the $(m_{1/2}, m_0)$ plane shown in panel (b) of Fig. 1, which includes the updated point K.

As mentioned earlier when discussing the specific benchmark points, the appropriate values of m_0 as functions of $m_{1/2}$ must be re-evaluated for the `ISASUGRA 7.67` code. A

similar minimization procedure to that discussed earlier has been repeated, parametrizing the shifts w.r.t. the values obtained with the **SSARD** code by a linear function of $m_{1/2}$, which have then been added to (5, 6, 8). The resulting parametrizations are:

$$\tan\beta = 5 : \quad m_0 = -9.96 + 0.197m_{1/2} + 4.02 \times 10^{-6}m_{1/2}^2, \quad (475 \leq m_{1/2} \leq 960) \quad (9)$$

$$\tan\beta = 10 : \quad m_0 = 13.57 + 0.142m_{1/2} + 4.90 \times 10^{-5}m_{1/2}^2, \quad (300 \leq m_{1/2} \leq 940) \quad (10)$$

$$\tan\beta = 20 : \quad m_0 = 50.39 + 0.142m_{1/2} + 6.77 \times 10^{-5}m_{1/2}^2, \quad (270 \leq m_{1/2} \leq 940), \quad (11)$$

$$\tan\beta = 35 : \quad m_0 = 89.56 + 0.239m_{1/2} + 5.70 \times 10^{-5}m_{1/2}^2, \quad (300 \leq m_{1/2} \leq 1050). \quad (12)$$

These have been used to produce the solid lines in Fig. 2.

We note in passing that, as the mass difference $\Delta M \equiv |m_{\tilde{\tau}_1} - m_\chi| \rightarrow 0$ towards the high- $m_{1/2}$ ends of the WMAP lines, these have portions at high $m_{1/2}$ where $\Delta M < m_\tau$. In these portions, the $\tilde{\tau}_1$ NLSP decays via a virtual τ , resulting in the dominance of four-body decay modes such as $\tilde{\tau}_1 \rightarrow \nu\chi\ell\bar{\nu}$ and $\nu\chi q\bar{q}$. In this case, the $\tilde{\tau}_1$ is stable on the scale of the size of the detector. This observation has important implications for $\tilde{\tau}_1$ observability at different colliders, as we discuss below.

3.2 Discussion of χ_2 Decay Branching Ratios

One of the key particles appearing in sparticle decay chains is the second neutralino χ_2 , whose branching ratios are quite model-dependent and have significant impact on sparticle detectability at future colliders. Moreover, χ_2 decays play crucial roles in reconstructing sparticle masses via cascade decays. Therefore, we now use **ISASUGRA 7.67** to discuss how the principal branching ratios of the χ_2 vary along the WMAP lines, noting several significant features that are important for phenomenology.

As seen in Fig 4(a) for $\tan\beta = 5$ and $\mu > 0$, χ_2 decays into $\tilde{\tau}\tau$ and other $\tilde{\ell}\ell$ modes dominate among the decay modes of interest. Both $\tilde{\tau}_1\tau$ and $\tilde{\tau}_2\tau$ contribute, the latter increasing with $m_{1/2}$ while the former decreases. The dip in the $\tilde{\ell}\ell$ branching ratio when $m_{1/2} \sim 300$ GeV reflects a similar switch between the $\tilde{\ell}_R\ell$ modes at low $m_{1/2}$ and the $\tilde{\ell}_L\ell$ modes at large $m_{1/2}$ ¹⁰. The most important other decay modes are the invisible $\chi_2 \rightarrow \tilde{\nu}\nu$ decays. The decay modes $\chi_2 \rightarrow \chi Z$ and $\chi_2 \rightarrow \chi h$ exhibit clear thresholds at $m_{1/2} \sim 270$ GeV and 310 GeV, respectively, reflecting the opening up of these two channels. We note that benchmark A is in the region of $m_{1/2}$ where the branching ratio for $\chi_2 \rightarrow \chi h$ is already declining from its peak value $\simeq 6$ %, while that for $\chi_2 \rightarrow \chi Z$ has already been reduced from its peak value $\simeq 0.6$ %.

¹⁰We note in passing that the branching ratio for $\tilde{\ell} \rightarrow \chi\ell$ exceeds 50 % for all the WMAP range of $m_{1/2}$.




Figure 4: Dominant branching ratios of the next-to-lightest neutralino χ_2 as functions of $m_{1/2}$ along the WMAP lines for (a) $\tan \beta = 5$, (b) $\tan \beta = 10$, (c) $\tan \beta = 20$, (d) $\tan \beta = 35$ and (e) $\tan \beta = 50$, all for $\mu > 0$, and (f) $\tan \beta = 10$ for $\mu < 0$. The locations of updated benchmark points along these WMAP lines are indicated, as are the upper limits on $m_{1/2}$ in panels (a, b, c) and (f).

Similar features are exhibited in Fig 4(b) for $\tan\beta = 10$ and $\mu > 0$, with the peak in the branching ratio for $\chi_2 \rightarrow \chi h$ reduced to $\simeq 5\%$ and that for $\chi_2 \rightarrow \chi Z$ increased to $\simeq 0.9\%$. The updated benchmark point B with $m_{1/2} = 250$ GeV is below both thresholds, whilst benchmark C has a near-maximal branching ratio for $\chi_2 \rightarrow \chi h$.

Panel (c) of Fig. 4 shows the corresponding χ_2 decay branching ratios for $\tan\beta = 20$ and $\mu > 0$. We see that the $\tilde{\tau}\tau$ mode is larger than for $\tan\beta = 10$, reflecting the larger τ Yukawa coupling. The dip in the $\tilde{\ell}\ell$ has moved to larger $m_{1/2} \sim 400$ GeV. The thresholds in $\chi_2 \rightarrow \chi Z$ and $\chi_2 \rightarrow \chi h$ are again prominent, with the former branching ratio now reaching slightly more than 1% . Benchmark G is located near the peak in $\chi_2 \rightarrow \chi h$ and the dip in $\chi_2 \rightarrow \tilde{\ell}\ell$, whilst benchmark H has almost equal branching ratios for $\tilde{\ell}\ell$ and $\tilde{\tau}\tau$, and smaller branching ratios for χh and particularly χZ .

When $\tan\beta = 35$ and $\mu > 0$, shown in panel (d) of Fig. 4, $\chi_2 \rightarrow \tilde{\tau}\tau$ becomes the dominant branching ratio for all values of $m_{1/2}$ allowed by WMAP. Apart from this, the most noticeable feature is the relative suppression of the $\chi_2 \rightarrow \tilde{\ell}\ell$ branching ratio, whose dip has now moved up to $m_{1/2} \sim 700$ GeV. The branching ratio for $\chi_2 \rightarrow \chi h$ exhibits a broad peak of similar height to the lower values of $\tan\beta$, whereas the branching ratio for $\chi_2 \rightarrow \chi Z$ is somewhat smaller. Benchmark I is located near the peaks of the $\chi_2 \rightarrow \chi h, \chi Z$ branching ratios, and benchmark J is located in the region of $m_{1/2}$ where the $\chi_2 \rightarrow \tilde{\ell}\ell$ starts to rise.

Panel (e) of Fig. 4 shows the branching ratios for $\tan\beta = 50$ and $\mu > 0$. Here we note the strong dominance of $\chi_2 \rightarrow \tilde{\tau}\tau$ decays, the approximate constancies (at relatively low levels) of the branching ratios for $\chi_2 \rightarrow \chi h, Z$ above their respective thresholds, and the strong suppression of $\chi_2 \rightarrow \tilde{\ell}\ell$ decays. Benchmark L has branching ratios that are typical for this value of $\tan\beta$.

Finally, panel (f) of Fig. 4 shows the branching ratios for $\tan\beta = 10$ when $\mu < 0$. In this case, there is dip structure in the branching ratio for $\chi_2 \rightarrow \tilde{\ell}\ell$, which rises monotonically with $m_{1/2}$, while the branching ratio for $\chi_2 \rightarrow \tilde{\tau}\tau$ is always large. In this case, the peak branching ratio for $\chi_2 \rightarrow \chi h$ is below that for $\chi_2 \rightarrow \chi Z$.

The χ_2 branching ratios are rather different at the tip of the funnel for $\tan\beta = 50$ - which has $\text{BR}(\chi_2 \rightarrow \chi h) = 0.11$ at the location of point M, by point K on the side of the rapid-annihilation funnel for $\tan\beta = 35$ and $\mu < 0$ - where the mode $\chi_2 \rightarrow \chi h$ gives half of the total χ_2 decay rate, and at the focus points E and F - where the χ_2 decay rate is saturated by $\chi q\bar{q}$ and by χh , respectively.

4 Production and Detectability along WMAP lines

We now study the reaches at different accelerators in the number of observable supersymmetric particles, and the changes in experimental signatures and topologies along the WMAP lines for different values of $\tan\beta$.

4.1 LHC

4.1.1 Sparticle Signatures

In order to visualize the changes in the signatures of sparticle decays along the WMAP lines, we first compute the mean numbers of different particle species produced in sparticle decay chains at LHC. Large samples of inclusive supersymmetric events have been generated along the WMAP lines, using PYTHIA 6.215 [34] interfaced to ISASUGRA 7.67 [20] to compute the average numbers of different particle species produced per supersymmetric event, and the production cross sections of exclusive and inclusive sparticle production reactions. Fig. 5 shows the average numbers of (a) Z , (b) h bosons, (c) τ leptons and (d) trilepton events with three ℓ ($\ell = e, \mu$) in inclusive sparticle decays at the LHC. The plots are all for $\mu > 0$ and display results for $\tan\beta = 10, 20, 35$ and 50 along the corresponding WMAP lines.

We see in panel (a) of Fig. 5 that the fractions of sparticle decays with a Z boson in the cascade is never large in the regions of the parameter space traversed by the WMAP lines, reaching about $0.1 Z^0/\text{sparticle decay}$ at large $m_{1/2}$ and $\tan\beta$.

The decays of sparticles into the lightest Higgs boson h , followed by $h \rightarrow b\bar{b}$ decay, provide another important signature at the LHC. These appear above the threshold for $\chi_2 \rightarrow \chi h$ decay at about $m_{1/2}=350$ GeV, and may reach 7-10% per sparticle event at larger $m_{1/2}$, as seen in panel (b) of Fig. 5.

As seen in panel (c), the numbers of τ leptons produced in sparticle events are always large, particularly at large $\tan\beta$ where there are between one and two per event. Finally, we see in panel (d) of Fig. 5 that the fraction of trilepton events is always above 10%, and may attain $\sim 40\%$ at large $m_{1/2}$ and small $\tan\beta$.

These plots reinforce the importance of the τ signature for sparticle detection at the LHC, within the CMSSM framework used here. Clearly the $\chi_2 \rightarrow \chi h, Z$ signatures are also interesting, but they may be quite challenging to exploit. However, we would like to emphasize that other decay patterns should not be neglected, and may be preferred in other supersymmetric models. For example, if the soft supersymmetry-breaking scalar masses for the Higgs multiplets are non-universal, $\chi_2 \rightarrow \chi h, Z$ decays may be more copious.




Figure 5: *Supersymmetric event signatures at the LHC: the numbers of Z bosons (upper left), light Higgs bosons h (upper right), τ leptons (lower left) and three-lepton final states per supersymmetric event are shown as functions of $m_{1/2}$ along the WMAP lines for four values of $\tan \beta$, all with $\mu > 0$. These plots were obtained with PYTHIA 6.215 [34] interfaced to ISASUGRA 7.67 [20].*

4.1.2 Detectability at the LHC

We now estimate the numbers of different species of supersymmetric particles that may be detectable at the LHC as functions of $m_{1/2}$ along the WMAP lines for different values of $\tan\beta$, using the following criteria.

- *Higgs bosons*: We generally follow the ATLAS and CMS studies of the numbers of observable bosons as functions of M_A and $\tan\beta$ [35]. In contrast to [3], here we also consider $H/A \rightarrow \chi_2\chi_2$ decays.

- *Gauginos*: Our criteria for the observability of heavier neutralinos at the LHC have been refined compared to our previous publication [3]. In particular, we first compute the total numbers of χ_2 produced in all sparticle events and the branching fraction corresponding to a dilepton final state. We consider this to be observable at the LHC if the product is at least 0.01 pb, corresponding to 1000 events produced with 100 fb^{-1} of integrated luminosity. The lightest neutralino χ is considered always to be detectable via the cascade decays of observed supersymmetric particles.

- *Gluinos*: These are considered to be observable for masses below 2.5 TeV [36].

- *Squarks*: The spartners of the lighter quark flavours u, d, s, c are considered to be observable if $m_{\tilde{q}} < 2.5 \text{ TeV}$ [36], but we recall that it is not known how to distinguish the spartners of different light-quark flavours at the LHC. In general, we assume that the stops and sbottoms $\tilde{t}, \tilde{b}$ are observable only if they weigh below 1 TeV, unless the gluino weighs $< 2.5 \text{ TeV}$ and the stop or sbottom can be produced in its two-body decays.

- *Charged Sleptons*: These are considered to be observable when the mass splitting $m_{\tilde{\ell}} - m_\chi > 30 \text{ GeV}$ ¹¹ and the inclusive production cross section, which includes their direct production and that in the decays of other supersymmetric particles, exceeds 0.1 pb, giving at least 10000 events with 100 fb^{-1} of integrated luminosity¹².

- *Sneutrinos*: These have not been considered as observable at the LHC, due to their large yield of invisible $\tilde{\nu} \rightarrow \nu\chi$ decays.

We first note the key differences between the expected numbers of detectable MSSM particles at the updated benchmark points, shown in Fig. 3, and the analogous LHC analysis in [3]. However, we would like to caution the reader that it is impossible to be precise about the capabilities of the LHC (or other accelerator) without detailed simulations that go beyond the scope of this paper.

¹¹This relatively conservative cut is to ensure that the $m_{\tilde{\ell}}$ decay lepton should be observable, and causes sleptons to be lost in some scenarios, particularly with the lower values of m_0 at the updated benchmark points.

¹²This criterion is illustrative: in a more detailed study, one should also include backgrounds such as W^+W^- , that may be very important.

- At point B, we now consider the heavier neutral Higgs bosons to be detectable via their decays into the χ_2 and into τ leptons, and the $H^\pm \rightarrow \tau\nu$ decays should also be detectable.

- We no longer consider the $H^\pm$ to be observable at point C.

- At point F, we now observe that the $\tilde{g}$ has branching ratios of 16 % and 17 %, respectively, for decays into $\chi_{2,3}\bar{t}t$, followed by branching ratios of 97 % and 99 % for $\chi_2 \rightarrow \chi h$ and $\chi_3 \rightarrow \chi Z$, respectively. The $\tilde{g}$ also has a branching ratio of 33 % for the decay into $\chi^\pm \bar{t}b$. We now consider that the $\chi_{2,3}$ should be detectable at this point, but not the $\chi^\pm$.

- At point H, supersymmetric particles are now well within the reach of the LHC, essentially because of the reduction in $m_{1/2}$. We now consider the $\tilde{q}$, $\tilde{g}$, χ and χ_2 to be detectable at this point.

- At point J, we now consider the χ_2 to be observable because of its large production rate, even though its decays generally include τ leptons.

- At point K, we now find that no sparticles are observable with our present criteria, although the lightest Higgs boson h is observable.

A corollary of these changes is that supersymmetric particles appear to be detectable at the LHC at all the updated benchmark points except K and M, which are located in rapid-annihilation funnels.

We now display in Fig. 6, as functions of $m_{1/2}$ along the WMAP lines for different values of $\tan\beta$, the numbers of different types of MSSM particles that should be observable at the LHC with 100 fb^{-1} of integrated luminosity in each of ATLAS and CMS. The nominal lower bounds on $m_{1/2}$ imposed by m_h (dashed lines) and $b \rightarrow s\gamma$ (dot-dashed lines) are also shown. These each have some uncertainties, for example **FeynHiggs** has a quoted error of $\sim 2 \text{ GeV}$ in the calculation of m_h so that (specifically) benchmark B in the top right panel should not be regarded as excluded.

We use the criteria explained above and generalize the results for the indicated benchmark scenarios that were shown in Fig. 3. In the top left panel, for $\tan\beta = 5$, we note first that only one MSSM Higgs boson is expected to be visible. In the top right panel for $\tan\beta = 10$, we see that the other Higgs bosons are also expected to be observable at low $m_{1/2}$ in the neighbourhood of point B, for example via decays into $\chi_2\chi_2$. All of the Higgs bosons are expected to be observable over the entire range of $m_{1/2}$ for $\tan\beta = 35, 50$ and $\mu > 0$, but not for $\tan\beta = 10$ and $\mu < 0$ (bottom right panel). All of the charginos and neutralinos are expected to be observable at low $m_{1/2}$, but only the lightest neutralino χ at high $m_{1/2}$. Since the sneutrinos decay invisibly and the rates for $\tilde{\ell}_2$ production are inadequate, only the $\tilde{\ell}_1$ are considered to be observable at the LHC for this value of $\tan\beta$, and only at low $m_{1/2}$. However, all the squarks and gluinos are expected to be observable anywhere along any of the

WMAP lines, except that for $\tan\beta = 50$ and $\mu > 0$, where sparticles would be unobservable at the end of the funnel, as exemplified by benchmark point M in Fig 3.

In the cases of larger values of $\tan\beta$, shown in the other panels of Fig. 6, all the MSSM Higgs bosons are expected to be observable for all the WMAP ranges of $m_{1/2}$. However, the observabilities of the different charginos and neutralinos resemble those for $\tan\beta = 10$. Some of the $\tilde{\ell}_2$ may be observable, at least for small $m_{1/2}$. As for $\tan\beta = 10$, all the squarks and gluinos are expected to be observable at the LHC anywhere along the parts of the WMAP lines that are shown, with the exception of the tip of the line for $\tan\beta = 50$.

We comment finally on the case of the metastable $\tilde{\tau}_1$ that appears towards the end of the WMAP lines where $\Delta M = |m_{\tilde{\tau}_1} - m_\chi| < m_\tau$. There is no specific study of this scenario at the LHC, but it has been estimated in an analogous gauge-mediated supersymmetry-breaking scenario that the efficiency for observing a metastable particle with $m = 640$ GeV would be 25 % (increasing for larger masses), and that a signal should be detectable if the overall production cross-section exceeds 1 fb [37]. In view of the large sparticle production cross sections at the LHC, and the fact (see Fig. 5) that each sparticle event produces on average 0.1 or more $\tilde{\tau}_1$ particles per event towards the ends of the WMAP lines, we believe that a metastable $\tilde{\tau}_1$ could be detected out to the ends of the coannihilation tails. On the other hand, it may not be easy to determine the mass accurately, as the peak measured from the relativistic $1/\beta$ factor broadens with increasing mass. For example, when $m = 640$ GeV the FWHM of the mass distribution is estimated to be about 250 GeV [37]. The error in the mass estimate may be reduced in a large statistical sample, but such an analysis would need to assume a realistic experimental environment in order to evaluate systematic errors.

4.2 Detectability at e^+e^- Linear Colliders

Our criteria for the observability of supersymmetric particles at linear colliders are based on their pair-production cross sections.

- Particles with cross sections in excess of 0.1 fb are considered as observable, thanks to their production in more than 100 events with an integrated luminosity of 1 ab^{-1} .
- The lightest neutralino χ is considered to be observable only through its production in the decays of heavier supersymmetric particles.
- Sneutrinos are considered to be detectable when the sum of the branching fractions for decays which lead to clean experimental signatures, such as $\tilde{\nu}_\ell \rightarrow \chi^\pm \ell^\mp$ ($\ell = e, \mu, \tau$) and $\tilde{\nu}_\tau \rightarrow W^+ \tilde{\tau}_1^-$, exceeds 15%.
- The $\gamma\gamma$ collider option at a linear collider would allow one to produce heavy neutral

Higgs bosons via the s -channel processes $\gamma\gamma \rightarrow A$ and $\gamma\gamma \rightarrow H$, extending the reach up to 750 GeV for 0.5 TeV $e^\pm$ beams and up to 1.5-2.0 TeV for 1.5 TeV $e^\pm$ beams. A $\gamma\gamma$ collider may also be used to look for gluinos, but we do not include this possibility in our analysis.

- Finally, we assume that a metastable $\tilde{\tau}_1$ could be detected at any linear e^+e^- collider with more than 100 events, and note that the mass could be measured more accurately than at the LHC, by measuring the production threshold as well as $1/\beta$.

We consider e^+e^- collision energies $\sqrt{s} = 0.5$ TeV, 1.0 TeV, 3 TeV and 5 TeV, and also the combined capabilities of the LHC and a 1-TeV linear collider. Comparing our present estimates of the physics reaches of linear e^+e^- colliders of different energies with those in [3], we observe changes due both to the criteria adopted and to the mass spectra. We note briefly the principal changes.

- The updated point F has no supersymmetric particles observable at 1 TeV centre-of-mass energy. This is in part due to differences in the ISAJET spectrum optimisation, where it now reproduces more closely that from SSARD.

- On the other hand, the reductions in the parameters $m_{1/2}, m_0$ for point H make slepton-pair production possible already below 1 TeV. The same point now has one squark accessible at CLIC with centre-of-mass energy 3 TeV, and all of the squarks at 5 TeV.

- Point M now has no squarks accessible even to CLIC operating with a centre-of-mass energy of 5 TeV.

4.2.1 TeV-Class e^+e^- Linear Colliders

The capabilities of a linear e^+e^- collider with $\sqrt{s} = 0.5$ TeV are illustrated in Fig. 7 [21]. It can see some gauginos and sleptons if $m_{1/2} \lesssim 500$ GeV. Its capabilities are therefore well suited to the $g_\mu - 2$ -friendlier scenarios displayed in the left columns of Fig. 3. Moreover, the sparticles that it would see would complement those detectable at the LHC. Also, we recall that such a linear collider would be able to measure sparticle properties much more accurately than the LHC [3, 8].

A linear e^+e^- collider able to deliver collisions at a centre of mass energy $\sqrt{s} = 1$ TeV has the potential to complement significantly the LHC in studying the supersymmetric spectrum, as seen in Fig. 8. In particular, such a linear collider would observe most sleptons and gauginos as long as $m_{1/2}$ is below about 600 GeV, as exemplified by benchmarks B, C, G, I and L. Also, it would detect at least one supersymmetric particle over the whole range of the WMAP lines, even up to their upper $m_{1/2}$ limits exemplified by benchmark point H, except in the funnel cases: $\mu > 0, \tan\beta = 50$ and $\mu < 0, \tan\beta = 35$. The accuracy it would provide in the measurements of several sparticle masses would enable GUT mass relations to

be tested, thereby completing the exploitation of the LHC data. We note also that a 1-TeV linear e^+e^- collider would be able to observe the lightest squark, the $\tilde{t}_1$, at low values of $m_{1/2}$. This possibility is exemplified by benchmark point B, as also seen in Fig. 3.

The complementarity between the LHC and 1-TeV linear e^+e^- collider is displayed clearly in Fig. 9, where we see that, together, they cover the majority of the MSSM spectrum over most of the $m_{1/2}$ ranges covered by the different WMAP lines.

4.2.2 CLIC

A 3-TeV lepton collider, such as CLIC, is expected to access almost all the sparticle spectrum for $m_{1/2} < 700$ GeV, as seen in Fig. 10. This would enable it, for example, to distinguish and provide detailed measurements of the different flavours of squarks that will not be possible at the LHC. Moreover, at larger $m_{1/2}$ CLIC would be able to observe (almost) all the spectrum of Higgs bosons, sleptons, charginos and neutralinos [23]. CLIC will also be able to observe gluinos via squark decays at the focus-point benchmark E¹³. It would therefore provide full complementarity with the LHC along the full extension of the lines.

Finally, we display in Fig. 11 the capabilities of a 5-TeV lepton collider such as CLIC. We see that all the MSSM particles are detected along all the WMAP lines, with the exception of the gluino that would generally have been seen at the LHC, and, for $\mu > 0$ and $\tan\beta = 50$, squarks towards the tip of the corresponding WMAP line, including the point M.

Despite the larger machine-induced backgrounds and beam energy spread, CLIC is expected to perform measurements of the properties of accessible supersymmetric particles with good accuracy [23]. Slepton and heavy Higgs boson masses can be determined to $\mathcal{O}(1\%)$ accuracy, also when accounting for realistic experimental conditions and resolutions. A similar accuracy can typically be obtained also for sparticles reconstructed through cascade decays, such as the χ_2 discussed below. The availability of polarized beams is not only beneficial to increase the signal cross sections (such as in the case of slepton-pair production) but also as an analyzing tool. A combination of measurements of the stop-pair production with different polarisation states can be used to determine the stop mixing angle [23].

4.3 A Neutralino Case Study

Many of the most interesting differences in the capabilities of the different colliders discussed above arise in the spectrum of charginos and neutralinos. In particular, while the LHC and a TeV-scale linear e^+e^- collider are largely complementary, they may not be able to discover

¹³As already noted, we do not discuss here the possibility of observing gluinos in $\gamma\gamma$ collisions.

all the charginos, neutralinos and sleptons when $m_{1/2} \gtrsim 400$ GeV, whereas CLIC can in principle observe all of them along all of the WMAP lines. To confirm these statements, we now compare the physics reaches of different accelerators for neutralinos in the $(m_{1/2}, m_0)$ plane, with the conclusions shown in Fig. 12(a).

For the LHC, we show Fig. 12(a) the region of the $(m_{1/2}, m_0)$ plane for $\tan \beta = 10$ in which dilepton structures due to the decay chains $\chi_2 \rightarrow \ell\tilde{\ell}, \tilde{\ell} \rightarrow \ell\chi$ and $\chi_2 \rightarrow XZ^0, Z^0 \rightarrow \ell^+\ell^-$ are expected to be observable in the cascade decays of heavier sparticles. We consider $\ell = e, \mu$ only, though the case $\ell = \tau$ would also be interesting at large $\tan \beta$, and require $\sigma \times$ dilepton branching ratio to exceed 0.01 pb, as discussed above. For our purposes, the relevant parts of the $(m_{1/2}, m_0)$ plane are those allowed by the WMAP dark matter constraint. We see that here the LHC coverage extends up to $m_{1/2} \simeq 500$ GeV, as also reflected in Fig. 6.

In order to see how accurately the 3-TeV version of CLIC could measure heavier neutralinos, we have made a new simulation of the process $e^+e^- \rightarrow \chi\chi_2$ followed by the decay chains $\chi_2 \rightarrow \ell\tilde{\ell}, \tilde{\ell} \rightarrow \ell\chi$ and $\chi_2 \rightarrow XZ^0, Z^0 \rightarrow \ell^+\ell^-$, using SIMDET [38] to simulate the detector response and PYTHIA 6.215 [34] interfaced to ISASUGRA [20] to simulate the signal. Events with at least two leptons and significant missing energy were selected. Both slepton and Standard Model gauge-boson pair-production backgrounds were considered. Combinatorial backgrounds were subtracted by taking the difference of the pair $e^+e^- + \mu^+\mu^-$ events and the mixed $e^\pm\mu^\mp$ events, and a sliding window has been used to search for a $>5\text{-}\sigma$ excess in the $M_{\ell\ell}$ mass distribution. The results shown in Figure 12(a) for $\tan \beta = 10$ make manifest the extended reach provided by CLIC, covering all the range of $m_{1/2}$ allowed by WMAP. At larger values of $\tan \beta$, decay chains involving the $\tilde{\tau}$ become more significant, requiring a more detailed study that should include τ reconstruction.

To benchmark the 3-TeV CLIC capabilities for measuring the masses of heavy neutralinos, a representative point has been chosen at $m_{1/2} = 700$ GeV, $m_0 = 150$ GeV, $\mu > 0$ and $\tan \beta = 10$, along the corresponding WMAP line. Here, $m_{\chi_2} = 540$ GeV, $m_\chi = 290$ GeV and $m_{\tilde{e}_L} = 490$ GeV. As seen in Fig. 12(b), at CLIC the dilepton invariant mass distribution shows a clear upper edge at 120 GeV due to $\chi_2 \rightarrow \ell^+\tilde{\ell}_L^-$ followed by $\tilde{\ell}_L^- \rightarrow \ell^-\chi$, which can be very accurately measured with 1 ab^{-1} of data. However, in order to extract the mass of the χ_2 state, the masses of both the $\tilde{\ell}_L$ and χ need to be known. In fact, making a two-parameter fit to the muon energy distribution, the masses of the $\tilde{\mu}_L$ and χ can be extracted with an accuracy of 3 % and 2.5 %, respectively, with 1 ab^{-1} of data [39]. Improved accuracy can be obtained with higher luminosity and using also a threshold energy scan. We therefore assume that these masses can be known to 1.7 % and 1.5 % respectively, which gives an uncertainty of 8 GeV, or 1.6 %, on the χ_2 mass, accounting for correlations. This uncertainty is dominated

by that in the $\tilde{\ell}$ and χ masses. The accuracy on the determination of the end point alone would correspond to a χ_2 mass determination to 0.3 GeV, fixing all other masses, which is not sensitive to the details of the beamstrahlung and accelerator-induced backgrounds.

5 Conclusions

We have seen in this paper how different colliders could provide complementary information about the MSSM spectrum in updated CMSSM benchmark scenarios and along lines in the $(m_{1/2}, m_0)$ planes for different values of $\tan\beta$ and both signs of μ . In addition to direct and indirect laboratory constraints, we have implemented the constraints on cold dark matter imposed by WMAP and other astrophysical and cosmological data.

We emphasize that the LSP is stable in any supersymmetric model in which R parity is conserved, such as the MSSM, in which case astrophysical and cosmological constraints on dark matter must be taken into account. In principle, the LSP might be some different sparticle, such as the gravitino or axino. Specific scenarios of this type include models with gauge-mediated supersymmetry breaking, and benchmarks for such models have been proposed elsewhere [8]. We recall, however, that the WMAP data on reionization of the early Universe disfavour models with warm dark matter [12], so that all such models must grapple with the constraint on cold dark matter provided by WMAP and other data.

Alternatively, one might seek to avoid the cold dark matter constraints by postulating some amount of R violation. The collider signatures of any such model would differ from those of the MSSM if R -violating couplings are not sufficiently small, and an additional set of astrophysical and cosmological constraints come into play if the R -violating couplings are not very large. Any R -violating scenario must address these issues.

We have updated a set of thirteen CMSSM benchmark points proposed previously [3] in light of the current WMAP constraint on supersymmetric dark matter. Since this constraint has reduced the dimensionality of the CMSSM parameter space, we have also introduced ‘WMAP lines’ in the $(m_{1/2}, m_0)$ plane for different values of $\tan\beta$, and have studied the physics reach of different accelerators, by charting the decay signatures and numbers of observable particles. It is interesting to observe that some important signatures, such as the pattern of χ_2 decays, remain rather uniform along these lines. However, we note that different signatures may be recovered by relaxing some of the assumptions taken here. An interesting example is the appearance of large χ_2 decay branching fractions into h^0 or Z^0 in models with non-universal Higgs masses. As these are of importance for defining the phenomenology of early runs at the LHC, we plan to consider them separately in conjunction with the

corresponding dark matter density constraints.

The observation at the LHC of an excess of events containing large missing transverse energy could immediately indicate that supersymmetry is present, and that R parity is conserved, so that the Universe should contain relic LSPs. However, it is possible that the relic LSP density might not saturate the bound on cold dark matter from WMAP, since there could be other significant contributions to the cold dark matter, *e.g.*, from axions or superheavy metastable particles. It is therefore interesting and important to study, within the CMSSM framework considered here, to what extent colliders can verify that the LSP ‘does its job’ of providing the cold dark matter [13], yielding a relic density that is neither too small nor too large.

The answer to this question requires a detailed study of the accuracy with which a given collider (or combination of colliders) can measure sparticle masses. General measurements of supersymmetric final states, based on the methods already established by ATLAS and CMS several years ago, involving, *e.g.*, the end-points of kinematic distributions, should allow the masses of the visible sparticles to be measured with precisions of 10 % or better in many cases. This should in turn provide constraints on the fundamental parameters of supersymmetry at the level of a few percent in minimal models such as the CMSSM. At this point one should be able to test the compatibility of the region of parameter space favoured by the LHC with that preferred by cold dark matter experiments and hypotheses, *e.g.*, with the WMAP lines.

To go further, one needs to understand the sensitivity of calculations of $\Omega_\chi h^2$ to variations in the supersymmetric model parameters [32]. These sensitivities depend on their values, are different for different benchmark scenarios, and vary along the WMAP lines. Determining in general the accuracy with which $\Omega_\chi h^2$ could be calculated would require detailed simulations going beyond the scope of this paper. However, existing studies of the previous version of benchmark point B yield a provisional answer in this case.

The following sparticle masses are thought to be measurable with the indicated accuracies at the LHC [40] at SPS point 1a [8], which is equivalent for our purposes to benchmark B:

$$\begin{aligned}
m_{\tilde{g}} &= 595.1(8.0), \\
m_{\tilde{q}_L} &= 540.3(8.8), \quad m_{\tilde{q}_L} = 520.4(11.8) \quad , \quad m_{\tilde{b}_1} = 491.9(7.5), \quad m_{\tilde{b}_2} = 524.5(7.9) \\
m_{\tilde{\ell}_L} &= 202.3(5.0), \quad m_{\tilde{\ell}_R} = 143.1(4.8), \quad m_{\tilde{\tau}_1} = 132.5(6.3), \\
m_\chi &= 96.2(4.8), \quad m_{\chi_2} = 176.9(4.7), \quad m_{\chi_4} = 377.9(5.1)
\end{aligned} \tag{13}$$

Since these simulations were made using ISASUGRA, we fit the determinations (13) to the CMSSM parameters $m_{1/2}, m_0$ and $\tan\beta$. We assume that the sign of μ is known, and set

$A_0 = 0$, recalling A_0 does not in any case have a large impact on $\Omega_\chi h^2$, and find:

$$m_0 = (103 \pm 8) \text{ GeV}, \quad m_{1/2} = (240 \pm 3) \text{ GeV}, \quad \tan \beta = 10.8 \pm 2. \quad (14)$$

We have adjusted the central value of m_0 to correspond to the updated benchmark value for point B, and then propagated the above uncertainties into the calculation of $\Omega_\chi h^2$ using **Micromegas**, with the result shown in Fig. 13.

As seen in Fig. 13, the calculated relic density has a well-behaved distribution that is well fit by a Gaussian with

$$\Omega_\chi h^2 = 0.111 \pm 0.018 \quad (15)$$

We have made a similar analysis using **SSARD**, finding an identical central value of $\Omega_\chi h^2$ and errors $^{+0.014}_{-0.020}$, $^{+0.001}_{-0.000}$ and $^{+0.010}_{-0.017}$ associated with m_0 , $m_{1/2}$ and $\tan \beta$, respectively. We conclude that collider measurements should, in principle enable the expected value of $\Omega_\chi h^2$ to be calculated with interesting accuracy. Measurements at linear e^+e^- colliders would increase the accuracy in (15) and also enable $\Omega_\chi h^2$ to be calculated for other benchmarks where data from the LHC alone might be insufficient.

The LHC and linear e^+e^- collider measurements would also provide crucial input for predicting the cross sections expected in direct dark matter searches or the LSP annihilation rate in the Sun. For this, it would in general be important to measure the higgsino and gaugino components of the LSP, which may be possible at the LHC in some regions of the parameter space by measuring the decay modes of the heavier gauginos or by comparing the rates of $\chi_2 \rightarrow \ell\ell\chi$ via a (virtual) slepton and $\chi_2 \rightarrow \chi h$.

Dark matter physics is another example of the complementarity of LHC and linear e^+e^- colliders for exploring new physics in the TeV energy range. Very possibly, Nature does not choose the CMSSM, and it is desirable to formulate benchmarks for other possibilities [8]. However, the CMSSM remains one of the prime options for physics beyond the Standard Model that might be accessible to the next generation of colliders. We hope that this paper contributes to a better understanding of what they might achieve, given the present constraints on supersymmetry from laboratory experiments and cosmology, as updated here using information from WMAP.

Acknowledgments

The work of K.A.O. was supported partly by DOE grant DE-FG02-94ER-40823. We thank G. Belanger, F. Boudjema and A. Pukhov for useful discussions about **Micromegas**.

References

- [1] L. Maiani, *Proceedings of the 1979 Gif-sur-Yvette Summer School On Particle Physics*, 1; G. 't Hooft, in *Recent Developments in Gauge Theories, Proceedings of the Nato Advanced Study Institute, Cargese, 1979*, eds. G. 't Hooft *et al.*, (Plenum Press, NY, 1980); E. Witten, *Phys. Lett. B* **105**, 267 (1981).
- [2] I. Hinchliffe, F. E. Paige, M. D. Shapiro, J. Soderqvist and W. Yao, *Phys. Rev.* **D55** (1997) 5520; S. Abdullin *et al.* [CMS Collaboration], hep-ph/9806366; S. Abdullin and F. Charles, *Nucl. Phys.* **B547** (1999) 60; TESLA Technical Design Report, DESY-01-011, Part III, *Physics at an e^+e^- Linear Collider* (March 2001).
- [3] M. Battaglia *et al.*, *Eur. Phys. J. C* **22**, 535 (2001) [arXiv:hep-ph/0106204].
- [4] Joint LEP 2 Supersymmetry Working Group, *Combined LEP Chargino Results, up to 208 GeV*, http://lepsusy.web.cern.ch/lepsusy/www/inos_moriond01/charginos_pub.html; *Combined LEP Selectron/Smuon/Stau Results, 183-208 GeV*, http://lepsusy.web.cern.ch/lepsusy/www/sleptons_summer02/slep_2002.html.
- [5] LEP Higgs Working Group for Higgs boson searches, OPAL Collaboration, ALEPH Collaboration, DELPHI Collaboration and L3 Collaboration, *Search for the Standard Model Higgs Boson at LEP*, CERN-EP/2003-011, arXiv:hep-ex/0306033.
- [6] M. S. Alam *et al.*, [CLEO Collaboration], *Phys. Rev. Lett.* **74**, 2885 (1995), as updated in S. Ahmed *et al.*, CLEO CONF 99-10; BELLE Collaboration, BELLE-CONF-0003, contribution to the 30th International conference on High-Energy Physics, Osaka, 2000. See also K. Abe *et al.*, [Belle Collaboration], arXiv:hep-ex/0107065; L. Lista [BaBar Collaboration], arXiv:hep-ex/0110010; C. Degrassi, P. Gambino and G. F. Giudice, *JHEP* **0012**, 009 (2000) [arXiv:hep-ph/0009337]; M. Carena, D. Garcia, U. Nierste and C. E. Wagner, *Phys. Lett. B* **499**, 141 (2001) [arXiv:hep-ph/0010003]; P. Gambino and M. Misiak, *Nucl. Phys. B* **611**, 338 (2001); D. A. Demir and K. A. Olive, *Phys. Rev. D* **65**, 034007 (2002) [arXiv:hep-ph/0107329]; T. Hurth, arXiv:hep-ph/0212304.
- [7] H. N. Brown *et al.* [Muon g-2 Collaboration], *Phys. Rev. Lett.* **86**, 2227 (2001) [arXiv:hep-ex/0102017]; G. W. Bennett *et al.* [Muon g-2 Collaboration], *Phys. Rev. Lett.* **89**, 101804 (2002) [Erratum-ibid. **89**, 1219903 (2002)] [arXiv:hep-ex/0208001]; M. Davier, S. Eidelman, A. Hocker and Z. Zhang, *Eur. Phys. J. C* **27**, 497 (2003)

- [arXiv:hep-ph/0208177]; see also K. Hagiwara, A. D. Martin, D. Nomura and T. Teubner, arXiv:hep-ph/0209187; F. Jegerlehner, unpublished, as reported in M. Krawczyk, arXiv:hep-ph/0208076.
- [8] B. C. Allanach *et al.*, *Proc. of the APS/DPF/DPB Summer Study on the Future of Particle Physics (Snowmass 2001)* ed. N. Graf, Eur. Phys. J. C **25**, 113 (2002) [eConf **C010630**, P125 (2001)] [arXiv:hep-ph/0202233].
- [9] M. Carena, S. Heinemeyer, C. E. Wagner and G. Weiglein, arXiv:hep-ph/9912223, Eur. Phys. J. C **26**, 601 (2003) [arXiv:hep-ph/0202167].
- [10] G. L. Kane, J. Lykken, S. Mrenna, B. D. Nelson, L. T. Wang and T. T. Wang, Phys. Rev. D **67**, 045008 (2003) [arXiv:hep-ph/0209061].
- [11] H. Baer, T. Krupovnickas and X. Tata, arXiv:hep-ph/0305325.
- [12] C. L. Bennett *et al.*, arXiv:astro-ph/0302207; D. N. Spergel *et al.*, arXiv:astro-ph/0302209; H. V. Peiris *et al.*, arXiv:astro-ph/0302225.
- [13] J. R. Ellis, J. S. Hagelin, D. V. Nanopoulos, K. A. Olive and M. Srednicki, Nucl. Phys. B **238**, 453 (1984); see also H. Goldberg, Phys. Rev. Lett. **50**, 1419 (1983).
- [14] See, for example: J. R. Ellis, T. Falk, G. Ganis, K. A. Olive and M. Srednicki, Phys. Lett. **B510**, 236 (2001) [arXiv:hep-ph/0102098]; L. Roszkowski, R. Ruiz de Austri and T. Nihei, JHEP **0108**, 204 (2001) [arXiv:hep-ph/0106334]; A. Djouadi, M. Drees and J. L. Kneur, JHEP **0108**, 055 (2001) [arXiv:hep-ph/0107316]; U. Chattopadhyay, A. Corsetti and P. Nath, Phys. Rev. D **66**, 035003 (2002) [arXiv:hep-ph/0201001]; H. Baer, C. Balazs and A. Belyaev, JHEP **0203**, 042 (2002) [arXiv:hep-ph/0202076]; R. Arnowitt and B. Dutta, arXiv:hep-ph/0211417; T. Kamon, R. Arnowitt, B. Dutta and V. Khotilovich, arXiv:hep-ph/0302249; H. Baer, C. Balazs, A. Belyaev, T. Krupovnickas and X. Tata, arXiv:hep-ph/0304303.
- [15] J. Ellis, K. A. Olive, Y. Santoso and V. C. Spanos, arXiv:hep-ph/0303043.
- [16] H. Baer and C. Balazs, arXiv:hep-ph/0303114; A. B. Lahanas and D. V. Nanopoulos, arXiv:hep-ph/0303130; U. Chattopadhyay, A. Corsetti and P. Nath, arXiv:hep-ph/0303201.
- [17] A. B. Lahanas, D. V. Nanopoulos and V. C. Spanos, Phys. Rev. D **62** (2000) 023515 [arXiv:hep-ph/9909497]; Mod. Phys. Lett. A **16** (2001) 1229 [arXiv:hep-ph/0009065];

- Phys. Lett. **B518** (2001) 94 [arXiv:hep-ph/0107151]; V. Barger and C. Kao, Phys. Lett. **B518** (2001) 117 [arXiv:hep-ph/0106189]; R. Arnowitt and B. Dutta, arXiv:hep-ph/0211417.
- [18] Information about this code is available from K. A. Olive: it contains important contributions from T. Falk, G. Ganis, J. McDonald, K. A. Olive and M. Srednicki.
- [19] S. Heinemeyer, W. Hollik and G. Weiglein, Comput. Phys. Commun. **124**, 76 (2000) [hep-ph/9812320] and Eur. Phys. J. C **9** (1999) 343 [arXiv:hep-ph/9812472].
- [20] We use mainly the 7.67 version of ISASUGRA, whose basic documentation is found in H. Baer, F. E. Paige, S. D. Protopopescu and X. Tata, *ISAJET 7.48: A Monte Carlo event generator for pp , $\bar{p}p$, and e^+e^- reactions*, hep-ph/0001086. The latest update is available from <http://paige.home.cern.ch/paige/>.
- [21] S. Matsumoto *et al.* [JLC Group], *JLC-1*, KEK Report 92-16 (1992); J. Bagger *et al.* [American Linear Collider Working Group], *The Case for a 500-GeV e^+e^- Linear Collider*, SLAC-PUB-8495, BNL-67545, FERMILAB-PUB-00-152, LBNL-46299, UCRL-ID-139524, LBL-46299, Jul 2000, arXiv:hep-ex/0007022; T. Abe *et al.* [American Linear Collider Working Group Collaboration], *Linear Collider Physics Resource Book for Snowmass 2001*, SLAC-570, arXiv:hep-ex/0106055, hep-ex/0106056, hep-ex/0106057 and hep-ex/0106058; TESLA Technical Design Report, DESY-01-011, Part III, *Physics at an e^+e^- Linear Collider* (March 2001).
- [22] R. W. Assmann *et al.* [CLIC Study Team], *A 3-TeV e^+e^- Linear Collider Based on CLIC Technology*, ed. G. Guignard, CERN 2000-08; for more information about this project, see: <http://ps-div.web.cern.ch/ps-div/CLIC/Welcome.html>.
- [23] CLIC Physics Study Group, <http://clicphysics.web.cern.ch/CLICphysics/> and Yellow Report in preparation.
- [24] M. Drees, Y. G. Kim, M. M. Nojiri, D. Toya, K. Hasuko and T. Kobayashi, Phys. Rev. D **63**, 035008 (2001) [arXiv:hep-ph/0007202].
- [25] J. Ellis, T. Falk and K. A. Olive, Phys. Lett. B **444**, 367 (1998) [arXiv:hep-ph/9810360]; J. Ellis, T. Falk, K. A. Olive and M. Srednicki, *Astropart. Phys.* **13**, 181 (2000) [arXiv:hep-ph/9905481]; M. E. Gómez, G. Lazarides and C. Pallis, Phys. Rev. D **61**, 123512 (2000) [arXiv:hep-ph/9907261], Phys. Lett. B **487**, 313 (2000) [arXiv:hep-ph/0004028] and Nucl. Phys. B **B638**, 165 (2002) [arXiv:hep-ph/0203131]; R. Arnowitt,

- B. Dutta and Y. Santoso, *Nucl. Phys. B* **606**, 59 (2001) [arXiv:hep-ph/0102181]; T. Nihei, L. Roszkowski and R. Ruiz de Austri, *JHEP* **0207**, 024 (2002) [arXiv:hep-ph/0206266].
- [26] M. Drees and M. M. Nojiri, *Phys. Rev. D* **47**, 376 (1993) [arXiv:hep-ph/9207234]; H. Baer and M. Brhlik, *Phys. Rev. D* **53**, 597 (1996) [arXiv:hep-ph/9508321] and *Phys. Rev. D* **57**, 567 (1998) [arXiv:hep-ph/9706509]; H. Baer, M. Brhlik, M. A. Diaz, J. Ferrandis, P. Mercadante, P. Quintana and X. Tata, *Phys. Rev. D* **63**, 015007 (2001) [arXiv:hep-ph/0005027]; A. B. Lahanas, D. V. Nanopoulos and V. C. Spanos, *Mod. Phys. Lett. A* **16**, 1229 (2001) [arXiv:hep-ph/0009065]; J. R. Ellis, T. Falk, G. Ganis, K. A. Olive and M. Srednicki in [14]; A. B. Lahanas and V. C. Spanos, *Eur. Phys. J. C* **23**, 185 (2002) [arXiv:hep-ph/0106345].
- [27] J. L. Feng, K. T. Matchev and T. Moroi, *Phys. Rev. Lett.* **84** (2000) 2322; J. L. Feng, K. T. Matchev and T. Moroi, *Phys. Rev.* **D61** (2000) 075005; J. L. Feng, K. T. Matchev and F. Wilczek, *Phys. Lett.* **B482** (2000) 388.
- [28] A. Djouadi, J.-L. Kneur and G. Moultaka, *SuSpect: a Fortran Code for the Supersymmetric and Higgs Particle Spectrum in the MSSM*, arXiv:hep-ph/0211331. The latest version is available from <http://www.lpm.univ-montp2.fr:7082/~kneur/>.
- [29] B. C. Allanach, S. Kraml and W. Porod, *JHEP* **0303**, 016 (2003) [arXiv:hep-ph/0302102]; see also the web pages <http://cern.ch/kraml/comparison/compare.html> for comparisons between different codes and http://wwwth.cern.ch/susycom/compare_vrsn.html for comparisons between different versions of ISASUGRA.
- [30] The RunII D0 top group, see <http://www-d0.fnal.gov/Run2Physics/top/conf.html>.
- [31] D. Pierce and A. Papadopoulos, *Phys. Rev.* **D50**, 565 (1994) [hep-ph/9312248] and *Nucl. Phys.* **B430**, 278 (1994) [hep-ph/9403240]; W. de Boer, R. Ehret and D. I. Kazakov, *Z. Phys. C* **67**, 647 (1995) [arXiv:hep-ph/9405342]; D. M. Pierce, J. A. Bagger, K. Matchev and R. Zhang, *Nucl. Phys.* **B491**, 3 (1997) [hep-ph/9606211].
- [32] J. R. Ellis, K. Enqvist, D. V. Nanopoulos and F. Zwirner, *Mod. Phys. Lett. A* **1**, 57 (1986); R. Barbieri and G. F. Giudice, *Nucl. Phys. B* **306**, 63 (1988); J. R. Ellis and K. A. Olive, *Phys. Lett. B* **514**, 114 (2001) [arXiv:hep-ph/0105004]; J. R. Ellis, K. A. Olive and Y. Santoso, *New J. Phys.* **4**, 32 (2002) [arXiv:hep-ph/0202110].

- [33] G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, arXiv:hep-ph/0112278.
- [34] T. Sjostrand, L. Lonnblad and S. Mrenna, arXiv:hep-ph/0108264.
- [35] ATLAS Collaboration, *ATLAS detector and physics performance Technical Design Report*, CERN/LHCC 99-14/15 (1999); CMS Collaboration, Technical Proposal, CERN/LHCC 94-38 (1994).
- [36] See [35], and the second and third papers in [2].
- [37] M. Kazana, G. Wrochna and P. Zalewski, CMS Conference Report 1999-019, contributed to the European Physical Society Conference on High-Energy Physics, Tampere, 1999.
- [38] M. Pohl and H. J. Schreiber, arXiv:hep-ex/0206009.
- [39] M. Battaglia and M. Gruwe, arXiv:hep-ph/0212140.
- [40] G. Polesello, private communication.




Figure 6: Estimates of the numbers of MSSM particles that may be detectable at the LHC as functions of $m_{1/2}$ along the WMAP lines for $\mu > 0$ and $\tan \beta = 5, 10, 20, 35$ and 50, and for $\mu < 0$ and $\tan \beta = 10$. The locations of updated benchmark points along these WMAP lines are indicated, as are the nominal lower bounds on $m_{1/2}$ imposed by m_h (dashed lines) and $b \rightarrow s\gamma$ (dot-dashed lines).




Figure 7: Estimates of the numbers of MSSM particles that may be detectable at a 0.5-TeV linear e^+e^- collider as functions of $m_{1/2}$ along the WMAP lines for $\mu > 0$ and $\tan \beta = 5, 10, 20, 35$ and 50, and for $\mu < 0$ and $\tan \beta = 10$. The locations of updated benchmark points along these WMAP lines are indicated, as are the nominal lower bounds on $m_{1/2}$ imposed by m_h (dashed lines) and $b \rightarrow s\gamma$ (dot-dashed lines).




Figure 8: Estimates of the numbers of MSSM particles that may be detectable at a 1-TeV linear e^+e^- collider as functions of $m_{1/2}$ along the WMAP lines for $\mu > 0$ and $\tan \beta = 5, 10, 20, 35$ and 50, and for $\mu < 0$ and $\tan \beta = 10$. The locations of updated benchmark points along these WMAP lines are indicated, as are the nominal lower bounds on $m_{1/2}$ imposed by m_h (dashed lines) and $b \rightarrow s\gamma$ (dot-dashed lines).




Figure 9: Estimates of the combined numbers of MSSM particles that may be detectable at a combination of the LHC and a 1-TeV linear e^+e^- collider as functions of $m_{1/2}$ along the WMAP lines for $\mu > 0$ and $\tan \beta = 5, 10, 20, 35$ and 50, and for $\mu < 0$ and $\tan \beta = 10$. The locations of updated benchmark points along these WMAP lines are indicated, as are the nominal lower bounds on $m_{1/2}$ imposed by m_h (dashed lines) and $b \rightarrow s\gamma$ (dot-dashed lines).




Figure 10: Estimates of the numbers of MSSM particles that may be detectable at the 3-TeV version of the CLIC linear e^+e^- collider as functions of $m_{1/2}$ along the WMAP lines for $\mu > 0$ and $\tan\beta = 5, 10, 20, 35$ and 50, and for $\mu < 0$ and $\tan\beta = 10$. The locations of updated benchmark points along these WMAP lines are indicated, as are the nominal lower bounds on $m_{1/2}$ imposed by m_h (dashed lines) and $b \rightarrow s\gamma$ (dot-dashed lines).




Figure 11: Estimates of the numbers of MSSM particles that may be detectable at the 5-TeV version of the CLIC linear e^+e^- collider as functions of $m_{1/2}$ along the WMAP lines for $\mu > 0$ and $\tan \beta = 5, 10, 20, 35$ and 50, and for $\mu < 0$ and $\tan \beta = 10$. The locations of updated benchmark points along these WMAP lines are indicated, as are the nominal lower bounds on $m_{1/2}$ imposed by m_h (dashed lines) and $b \rightarrow s\gamma$ (dot-dashed lines).




Figure 12: (a) Comparison of the sensitivities in the $(m_{1/2}, m_0)$ plane of the LHC and CLIC at 3 TeV to $\chi_2 \rightarrow \chi \ell^+ \ell^-$ decay and (b) the dilepton mass spectrum from this decay observable at 3-TeV CLIC for the CMSSM point $m_{1/2} = 700$ GeV, $m_0 = 150$ GeV, $\mu > 0$ and $\tan \beta = 10$ discussed in the text.




Figure 13: *The estimated accuracy with which $\Omega_\chi h^2$ could be predicted on the basis of LHC data for the updated benchmark point B, using the projected experimental errors reported in [40], a fit to ISASUGRA 7.67 parameters and the Micromegas code. We find similar results using SSARD.*